

SDP Gaps from Pairwise Independence

Siavosh Benabbas

Konstantinos Georgiou

Avner Magen*

Madhur Tulsiani†

Received: April 14, 2010; revised: November 8, 2011; published: June 21, 2012.

Abstract: We consider the problem of approximating fixed-predicate constraint satisfaction problems (MAX k -CSP $_q(P)$), where the variables take values from $[q] = \{0, 1, \dots, q-1\}$, and each constraint is on k variables and is defined by a fixed k -ary predicate P . Familiar problems like MAX 3-SAT and MAX-CUT belong to this category. Austrin and Mossel recently identified a general class of predicates P for which MAX k -CSP $_q(P)$ is hard to approximate. They study predicates $P : [q]^k \rightarrow \{0, 1\}$ such that the set of assignments accepted by P contains the support of a balanced pairwise independent distribution over the domain of the inputs. We refer to such predicates as *promising*. Austrin and Mossel show that for any promising predicate P , the problem MAX k -CSP $_q(P)$ is Unique-Games-hard to approximate better than the trivial approximation obtained by a random assignment.

We give an unconditional analogue of this result in a restricted model of computation. We consider the hierarchy of semidefinite relaxations of MAX k -CSP $_q(P)$ obtained by augmenting the canonical semidefinite relaxation with the Sherali-Adams hierarchy. We show that for any promising predicate P , the integrality gap remains the same as the approximation ratio achieved by a random assignment, even after $\Omega(n)$ levels of this hierarchy.

*Funded in part by NSERC.

†Supported by the NSF grant CCF-083279.

ACM Classification: F.2.2, F.1.3, G.1.6, F.2.3

AMS Classification: 68Q17, 68W25, 68W20, 90C59

Key words and phrases: semidefinite and linear programming hierarchies, integrality gaps, constraint satisfaction

We also introduce a new general set of techniques to define consistent “local distributions” over partial assignments to variables in the problem. Defining such distributions has often been the crux of proving lower bounds on integrality gaps for hierarchies like the ones we consider.

1 Introduction

A constraint satisfaction problem (CSP) is defined by a set of constraints on a set of q -valued variables. In the maximization version (MAX-CSP) one tries to maximize the number of constraints that can be simultaneously satisfied. In this paper, we consider the most commonly studied families of CSPs, denoted by MAX k -CSP $_q(P)$, defined as follows. The variables take values over the fixed alphabet $[q] = \{0, 1, \dots, q-1\}$. All constraints are defined by a single predicate $P : [q]^k \rightarrow \{0, 1\}$ and have the form $P(x_1 + b_1 \bmod q, \dots, x_k + b_k \bmod q)$ for $b_i \in [q]$. For instance, MAX 3-SAT is the same as MAX 3-CSP $_2(\text{OR}_3)$, where OR_3 denotes the 3-variable OR function. We refer to the terms $x_i + b_i \bmod q$ as *literals*,¹ generalizing from the Boolean case, where the literals are x_i and $\neg x_i = x_i + 1 \bmod 2$.

Given a predicate P , an instance of the MAX k -CSP $_q(P)$ problem is a collection of constraints as above and the objective is to maximize the number of constraints that can be satisfied simultaneously. As special cases, we obtain many well-studied MAX-CSP problems, e. g., MAX k -SAT, MAX k -XOR, MAX k -LIN $_q$ (systems of mod q linear equations with k variables per equation), etc. Note that MAX 2-LIN $_2$ includes MAX-CUT. We use MAX k -CSP $_q$ to denote the union of the classes MAX k -CSP $_q(P)$ over all $P : [q]^k \rightarrow \{0, 1\}$.

The MAX k -CSP $_q$ problem is NP-hard for any $k \geq 2$, $q \geq 2$. A lot of effort has been devoted to determining the true approximability threshold of the problem as a function of k and q . For the Boolean case ($q = 2$), Samorodnitsky and Trevisan [25] proved that the problem is hard to approximate within a factor less than $2^k/2^{2\sqrt{k}}$, which was improved to $2^k/2^{\sqrt{2k}}$ by Engebretsen and Holmerin [12]. Later Samorodnitsky and Trevisan [24] showed that it is Unique-Games-hard (UG-hard) to approximate the same problem within a factor less than $2^k/2^{\lceil \log k + 1 \rceil}$. For the general case of q -valued variables (MAX k -CSP $_q$), Guruswami and Raghavendra [16] proved UG-hardness of approximation within a factor less than $q^k/(kq^2)$ when q is a prime.

Austrin and Mossel [5] studied the complexity of MAX k -CSP $_q(P)$ for predicates $P : [q]^k \rightarrow \{0, 1\}$ such that the set of accepted inputs $P^{-1}(1)$ contains the support of a balanced pairwise independent distribution on $[q]^k$. We shall refer to such predicates as *promising*. In a very general result, which (assuming the Unique Games Conjecture) subsumes all the above, Austrin and Mossel [5] showed that for a promising predicate P , the MAX k -CSP $_q(P)$ problem is UG-hard to approximate within a factor less than $q^k/|P^{-1}(1)|$. Considering that a random assignment satisfies a $|P^{-1}(1)|/q^k$ fraction of all the constraints, this is the strongest result one can get for such P . Using appropriate choices for the predicate P , this then implies a hardness ratio of $q^k/((1+o(1))kq^2)$ for MAX k -CSP $_q$ for any $q \geq 2$, a ratio of $q^k/(kq(q-1))$ when q is a prime power, and $2^k/(k+O(k^{0.525}))$ for $q = 2$.

In this paper, we study the inapproximability of MAX k -CSP $_q(P)$ for promising predicates P , in a restricted model of computation. The model we consider is the hierarchy of semidefinite relaxations

¹We believe this terminology was introduced by [5] (see also [4]).

of $\text{MAX } k\text{-CSP}_q(P)$, given by an application of the Sherali-Adams [29] strengthenings of the canonical semidefinite relaxation of $\text{MAX } k\text{-CSP}_q(P)$. We give an unconditional analogue of the result of Austrin and Mossel in this hierarchy, demonstrating that even after $\Omega(n)$ levels of this hierarchy, the integrality gap remains at least $q^k/|P^{-1}|$, which is the approximation ratio achieved by a random assignment. (The implied constant in the $\Omega(n)$ notation depends on k and q .)

Hierarchies of linear and semidefinite programs

A standard approach in approximating NP-hard problems, and therefore $\text{MAX } k\text{-CSP}_q$, is to formulate the problem as a 0-1 integer program and then relax the integrality condition to get a linear (or semidefinite) program which can be solved efficiently. The quality of such an approach is intimately related to the *integrality gap* of the relaxation, namely, the ratio between the optimum of the relaxation and that of the integer program.

Several methods (or procedures) were developed in order to obtain tightenings of relaxations in a systematic manner. These procedures give a sequence or a *hierarchy* of increasingly tighter relaxations of the starting program. The commonly studied ones include the hierarchies defined by Lovász-Schrijver [19], Sherali-Adams [29], and Lasserre [17] (see [18] for a comparison). Stronger relaxations in the sequence are referred to as higher *levels* of the hierarchy. It is known for all these hierarchies that for a starting program with n variables, the program at level n has integrality gap 1, and that it is possible to optimize over the program at the r th level in time $n^{O(r)}$.

Many known linear (semidefinite) programs can be captured by constant many levels of the Sherali-Adams (Lasserre) hierarchy. In fact, these semidefinite programs can also be captured by a “mixed” hierarchy, first studied by Raghavendra² [22]; where we augment the basic semidefinite relaxation by adding new real variables and imposing linear constraints according to the Sherali-Adams hierarchy. We will refer to this hierarchy of programs as the *Sherali-Adams SDP hierarchy*.

Fernandez de la Vega and Kenyon-Mathieu [14] have provided a PTAS for MAX-CUT in dense graphs using Sherali-Adams LP hierarchy. In [20] it is shown how to get a Sherali-Adams based PTAS for Vertex-Cover and Max-Independent-Set in minor-free graphs, while recently Mathieu and Sinclair [21] showed that the integrality gap for the matching polytope is asymptotically $1 + 1/r$, and Bateni, Charikar and Guruswami [6] that the integrality gap for a natural LP formulation of the MaxMin allocation problem is at most $n^{1/r}$, both after r many Sherali-Adams tightenings. Chlamtac [10] and Chlamtac and Singh [11] gave an approximation algorithm for Max-Independent-Set in hypergraphs based on the Lasserre hierarchy, with the performance depending on the number of levels. Recently, an $O(n^{1/4})$ approximation for Densest k -Subgraph was shown by Bhaskara et al. [7], using linear programs implied by $O(\log n)$ levels of the Lovász-Schrijver hierarchy.

Lower bounds in these hierarchies amount to showing that the integrality gap remains large even after many levels of the hierarchy. Integrality gaps for $\Omega(n)$ levels can be seen as unconditional lower bounds (as they rule out even exponential time algorithms obtained by the hierarchy) in a restricted (but still fairly interesting) model of computation. Considerable effort has been invested in proving such lower bounds (see [3, 31, 30, 28, 8, 13, 1, 2, 27, 15, 14]). For some CSPs in particular, strong lower bounds ($\Omega(n)$ levels) have recently been proved for the Lasserre hierarchy (which is the strongest) by [26] and [32],

²The hierarchies studied by Raghavendra were in fact slightly weaker than the ones defined here.

who showed a factor 2 integrality gap for MAX k -XOR and factor $2^k/(2k)$ integrality gap for MAX k -CSP respectively.

In a beautiful result, Raghavendra [22] showed a general connection between integrality gaps and UG-hardness results. His result essentially shows that for MAX k -CSP $_q(P)$, if the integrality gap of a program obtained by k levels of the Sherali-Adams SDP hierarchy is I , then the MAX k -CSP $_q(P)$ is UG-hard to approximate better than a factor of I . However, in our case the hardness is already known (by the work of Austrin and Mossel), and we are interested in finding the integrality gap for programs obtained by $\Omega(n)$ levels.

Our result and techniques

Both the known results in the Lasserre hierarchy (and previous analogues in the Lovász-Schrijver hierarchy) seemed to heavily rely on the structure of the predicate for which the integrality gap was proved; in particular, the predicate is always some system of linear equations. It was not clear if the techniques could be extended using only the fact that the predicate is promising (which is a much weaker condition). In this paper, we try to explore this issue, proving $\Omega(n)$ level gaps for the Sherali-Adams SDP hierarchy for any promising predicate.

Theorem 1.1. *Let $P : [q]^k \rightarrow \{0, 1\}$ be a predicate such that $P^{-1}(1)$ contains the support of a balanced pairwise independent distribution μ . Then for every constant $\zeta > 0$, there exist $c = c(q, k, \zeta) > 0$ such that for large enough n , the integrality gap of MAX k -CSP $_q(P)$ for the tightening obtained by cn levels of the Sherali-Adams SDP hierarchy³ is at least*

$$\frac{q^k}{|P^{-1}(1)|} - \zeta.$$

Remark 1.2. We note that weaker integrality gaps for these predicates also follow, via reductions, from the corresponding integrality gap results for Unique Games. In particular, a $(\log \log n)^{\Omega(1)}$ -level gap for the SDP hierarchy discussed above, follows from the recent results of Raghavendra and Steurer [23]. Also, $\Omega(n^\delta)$ -level gaps (where $\delta \rightarrow 0$ as $\zeta \rightarrow 0$) for the Sherali-Adams LP hierarchy can be deduced from the results of Charikar, Makarychev, and Makarychev [9].

A first step in achieving our result is to reduce the problem of a level- t gap to a question about a family of distributions over assignments associated with sets of variables of size at most t . These distributions should be (a) supported only on satisfying (partial) assignments, (b) should be consistent among themselves, in the sense that for $S_1 \subseteq S_2$ which are subsets of variables, the distributions over S_1 and S_2 should be equal on S_1 , and (c) should be balanced and pairwise-independent. The first requirement guarantees that the solution achieves objective value that corresponds to satisfying *all* the constraints of the instance. The second requirement implies feasibility for the Sherali-Adams LP constraints, while the last one makes it easy to produce vectors satisfying the semidefinite constraints.

The second step is to come up with these distributions! We explain why the simple method of picking a uniform distribution (or a reweighting of it according to the pairwise independent distribution that is supported by P) over the satisfying assignments cannot work. Instead we introduce the notion of “advice

³See the resulting SDP in Section 2.3.

sets.” These are sets on which it is “safe” to define such simple distributions. The actual distribution for a set S we use is then the one induced on S by a simple distribution defined on the advice-set of S . Getting such advice sets heavily relies on notions of expansion of the constraints graph. In particular, we use the fact that random instances have inherently good expansion properties. At the same time, such instances are highly unsatisfiable, ensuring that the resulting integrality gap is large.

Arguing that it is indeed “safe” to use simple distributions over the advice sets relies on the fact that the predicate P in question is promising, namely $P^{-1}(1)$ contains the support of a balanced pairwise independent distribution. We find it interesting and somewhat curious that the condition of pairwise independence comes up in this context for a reason very different than in the case of UG-hardness. Here, it represents the limit to which the expansion properties of a random CSP instance can be pushed to define such distributions.

2 Preliminaries and notation

2.1 Constraint satisfaction problems

For an instance Φ of MAX k -CSP $_q$, we denote the variables by $\{x_1, \dots, x_n\}$, their domain $\{0, \dots, q-1\}$ by $[q]$, and the constraints by C_1, \dots, C_m . Each constraint is a function of the form $C_i : [q]^{T_i} \rightarrow \{0, 1\}$ depending only on the values of the variables in the ordered tuple T_i with $|T_i| \leq k$.

For a set $S \subseteq [n]$ of variables, we denote by $[q]^S$ the set of all mappings from the set S to $[q]$. In the context of variables, these mappings can be understood as partial assignments to a given subset of variables. For $\alpha \in [q]^S$, we denote its projection to $S' \subseteq S$ by $\alpha(S')$. Also, for $\alpha_1 \in [q]^{S_1}$, $\alpha_2 \in [q]^{S_2}$ such that $S_1 \cap S_2 = \emptyset$, we denote by $\alpha_1 \circ \alpha_2$ the assignment over $S_1 \cup S_2$ defined by α_1 and α_2 .

We shall prove results for constraint satisfaction problems where every constraint is specified by the same Boolean predicate $P : [q]^k \rightarrow \{0, 1\}$. We denote the set of assignments for which the predicate evaluates to 1 by $P^{-1}(1)$. A CSP instance for such a problem is a collection of constraints of the form of P applied to k -tuples of *literals*. For a variable x with domain $[q]$, we take a literal to be $(x + a) \bmod q$ for any $a \in [q]$. We record this more formally in the following definition.

Definition 2.1. For a given $P : [q]^k \rightarrow \{0, 1\}$, an instance Φ of MAX k -CSP $_q(P)$ is a set of constraints C_1, \dots, C_m where each constraint C_i is over a k -tuple $T_i = \{x_{i_1}, \dots, x_{i_k}\}$ of variables and is of the form $P(x_{i_1} + a_{i_1}, \dots, x_{i_k} + a_{i_k})$ for some $a_{i_1}, \dots, a_{i_k} \in [q]$. We denote the maximum number of constraints that can be simultaneously satisfied by $\text{OPT}(\Phi)$.

2.2 Expanding CSP instances

For an instance Φ of MAX k -CSP $_q$, define its constraint graph G_Φ as the following bipartite graph from L to R . The left hand side L consists of a vertex for each constraint C_i . The right hand side R consists of a vertex for each variable x_j . There is an edge between a constraint-vertex i and a variable-vertex j , whenever variable x_j appears in constraint C_i . When Φ is clear from the context, we will abbreviate G_Φ by G .

For $C_i \in L$ we denote by $\Gamma(C_i) \subseteq R$ the set of neighbors of C_i in R . For a set $\mathcal{C} \subseteq L$ of constraints, $\Gamma(\mathcal{C})$ denotes $\bigcup_{C_i \in \mathcal{C}} \Gamma(C_i)$. For $S \subseteq R$, we call a constraint, $C_i \in L$, S -dominated if $\Gamma(C_i) \subseteq S$. We denote

maximize	$\sum_{i=1}^m \sum_{\alpha \in [q]^{T_i}} C_i(\alpha) \cdot X_{(T_i, \alpha)}$	
subject to	$\langle \mathbf{v}_{(i_1, j_1)}, \mathbf{v}_{(i_2, j_2)} \rangle = X_{(\{i_1, i_2\}, (j_1, j_2))}$	$\forall i_1 \neq i_2 \in [n], j_1, j_2 \in [q]$
	$\langle \mathbf{v}_{(i, j_1)}, \mathbf{v}_{(i, j_2)} \rangle = 0$	$\forall i \in [n], j_1 \neq j_2 \in [q]$
	$\ \mathbf{v}_{(i, j)}\ ^2 = \langle \mathbf{v}_{(i, j)}, \mathbf{v}_0 \rangle = X_{(\{i\}, j)}$	$\forall i \in [n], j \in [q]$
	$\ \mathbf{v}_0\ ^2 = X_{(\emptyset, \emptyset)} = 1$	
	$\sum_{j \in [q]} X_{(S \cup \{i\}, \alpha \circ j)} = X_{(S, \alpha)}$	$\forall S \text{ s. t. } S < t, \forall i \notin S, \alpha \in [q]^S$
	$X_{(S, \alpha)} \geq 0$	$\forall S \text{ s. t. } S \leq t, \forall \alpha \in [q]^S$

 Figure 1: SDP for MAX k -CSP $_q$ augmented with level- t Sherali-Adams constraints

by $G|_{-S}$ the bipartite subgraph of G that we get after removing S and all S -dominated constraints. Finally, we also denote by $\mathcal{C}(S)$ the set of all S -dominated constraints. Similarly, for $u \in R$ we denote by $\Gamma(u) \subseteq L$ the set of neighbors of u in L .

Our result relies on the expansion of the support sets of the constraints. We make this notion formal below.

Definition 2.2. Consider a bipartite graph $G = (V, E)$ with partition L, R . The *boundary expansion* of $X \subseteq L$ is the value $|\partial X|/|X|$, where $\partial X = \{u \in R : |\Gamma(u) \cap X| = 1\}$. G is (r, e) -*boundary expanding* if the boundary expansion for all (nonempty) subsets of L of size at most r is at least e .

2.3 The Sherali-Adams SDP hierarchy

Below we present a relaxation for the MAX k -CSP $_q$ problem as it is obtained by applying a level- t Sherali-Adams tightening to the basic SDP formulation of some instance Φ of MAX k -CSP $_q$. A well known fact states that the level- n Sherali-Adams tightening (even starting from a linear program) provides a perfect formulation, i. e., the integrality gap is 1 (see [29] or [18] for a proof).

The intuition behind the level- t Sherali-Adams tightening is the following. Note that an integer solution to the problem can be given by a single mapping $\alpha_0 \in [q]^{[n]}$, which is an assignment to all the variables. Using this, we can define 0/1 variables $X_{(S, \alpha)}$ for each $S \subseteq [n]$ such that $|S| \leq t$ and $\alpha \in [q]^S$. The intended solution is $X_{(S, \alpha)} = 1$ if $\alpha_0(S) = \alpha$ and 0 otherwise. We introduce $X_{(\emptyset, \emptyset)}$ which is intended to be 1. By relaxing the integrality constraint on the variables, we obtain the LP conditions given by level- t of the Sherali-Adams hierarchy.

We can further strengthen the integer program by adding the quadratic constraints

$$X_{(\{i_1, i_2\}, (j_1, j_2))} = X_{(\{i_1\}, j_1)} \cdot X_{(\{i_2\}, j_2)}$$

for $i_1, i_2 \in [n]$ and $j_1, j_2 \in [q]$.⁴ As solving quadratic programs is NP-hard we then relax these quadratic

⁴Abusing notation slightly, here we use $X_{(\{i_1, i_2\}, (j_1, j_2))}$ to denote $X_{(\{i_1, i_2\}, \alpha)}$ where α is the function mapping i_1 to j_1 and i_2 to j_2 .

constraints to the existence of vectors $\mathbf{v}_{(i,j)}$ and a unit vector \mathbf{v}_0 , for which we require that

$$\langle \mathbf{v}_{(i_1,j_1)}, \mathbf{v}_{(i_2,j_2)} \rangle = X_{(\{i_1,i_2\},\{j_1,j_2\})} \quad \text{and} \quad \langle \mathbf{v}_{(i,j)}, \mathbf{v}_0 \rangle = X_{(\{i\},j)}.$$

The complete relaxation can be seen in [Figure 1](#).

For an SDP formulation of MAX k -CSP $_q$, and for a given instance Φ of the problem, we denote by $\text{FRAC}(\Phi)$ the SDP (fractional) optimum, and by $\text{OPT}(\Phi)$ the integral optimum. For the particular instance Φ , the integrality gap is then defined as $\text{FRAC}(\Phi)/\text{OPT}(\Phi)$. The integrality gap of the formulation is the supremum of integrality gaps over all instances.

Next we give a sufficient condition for the existence of a solution to the program obtained by level- t of the Sherali-Adams SDP hierarchy for a MAX k -CSP $_q$ instance Φ .

Lemma 2.3. *Consider a family $\{\mathcal{D}(S)\}_{S \subseteq [n]: |S| \leq t}$ of distributions, where each $\mathcal{D}(S)$ is defined over $[q]^S$. Suppose that for every $S \subseteq T \subseteq [n]$ with $|T| \leq t$, the distributions $\mathcal{D}(S), \mathcal{D}(T)$ are equal on S and there exists a set $\{\mathbf{v}_{(i,j)}\}_{i \in [n], j \in [q]}$ of vectors and a unit vector \mathbf{v}_0 satisfying:*

1. for all $i \in [n]$ and $j_1 \neq j_2$, $\langle \mathbf{v}_{(i,j_1)}, \mathbf{v}_{(i,j_2)} \rangle = 0$,
2. for all $i \in [n], j \in [q]$, $\langle \mathbf{v}_{(i,j)}, \mathbf{v}_0 \rangle = \|\mathbf{v}_{(i,j)}\|^2 = \Pr_{\mathcal{D}(\{i\})}[j]$, and
3. for all $i_1 \neq i_2 \in [n]$ and $j_1, j_2 \in [q]$, $\langle \mathbf{v}_{(i_1,j_1)}, \mathbf{v}_{(i_2,j_2)} \rangle = \Pr_{\mathcal{D}(\{i_1,i_2\})}[(j_1, j_2)]$.

Then the vectors together with the LP variables $X_{(S,\alpha)} = \Pr_{\mathcal{D}(S)}[\alpha]$ form a feasible solution to the program in [Figure 1](#).

Proof. Consider some $S \subseteq [n]$, $|S| < t$, and some $i \notin S$. Note that the distributions $\mathcal{D}(S), \mathcal{D}(S \cup \{i\})$ are equal on S , and therefore we have

$$\begin{aligned} \sum_{j \in [q]} X_{(S \cup \{i\}, \alpha \circ j)} &= \sum_{j \in [q]} \Pr_{\beta \sim \mathcal{D}(S \cup \{i\})}[\beta = \alpha \circ j] \\ &= \sum_{j \in [q]} \Pr_{\beta \sim \mathcal{D}(S \cup \{i\})}[(\beta(i) = j) \wedge (\beta(S) = \alpha)] \\ &= \Pr_{\beta \sim \mathcal{D}(S \cup \{i\})}[\beta(S) = \alpha] \\ &= \Pr_{\beta' \sim \mathcal{D}(S)}[\beta' = \alpha] \\ &= X_{(S,\alpha)}. \end{aligned}$$

The same argument for $S = \emptyset$ shows that $X_{(\emptyset,\emptyset)} = 1$. It is clear that the solution satisfies all the other required conditions by definition, which proves the lemma. \square

2.4 Pairwise independence and approximation resistant predicates

We say that a distribution μ over variables x_1, \dots, x_k , is a balanced pairwise independent distribution over $[q]^k$, if we have

$$\forall j \in [q]. \forall i. \Pr_{\mu}[x_i = j] = \frac{1}{q} \quad \text{and} \quad \forall j_1, j_2 \in [q]. \forall i_1 \neq i_2. \Pr_{\mu}[(x_{i_1} = j_1) \wedge (x_{i_2} = j_2)] = \frac{1}{q^2}.$$

A predicate P is called approximation resistant if it is hard to approximate the $\text{MAX } k\text{-CSP}_q(P)$ problem better than using a random assignment. Assuming the Unique Games Conjecture, Austrin and Mossel [5] show that a predicate is approximation resistant if it is possible to define a balanced pairwise independent distribution μ such that P is always 1 on the support of μ .

Definition 2.4. A predicate $P : [q]^k \rightarrow \{0, 1\}$ is called *promising* if there exist a distribution supported over a subset of $P^{-1}(1)$ that is pairwise independent and balanced. If μ is such a distribution we say that P is promising supported by μ .

3 Towards defining consistent distributions

To construct valid solutions for the Sherali-Adams SDP hierarchy, we need to define distributions over every set S of bounded size as is required by Lemma 2.3. Since we will deal with promising predicates supported by some distribution μ , in order to satisfy consistency between distributions we will heavily rely on the fact that μ is a balanced pairwise independent distribution.

Assume for simplicity that μ is uniform over $P^{-1}(1)$ (the intuition for the general case is not significantly different). It is instructive to think of $q = 2$ and the predicate P being $k\text{-XOR}$, $k \geq 3$. Observe that the uniform distribution over $P^{-1}(1)$ is pairwise independent and balanced. A first attempt would be to define for every S , the distribution $\mathcal{D}(S)$ as the uniform distribution over all consistent assignments of S . We argue that such distributions are in general problematic. This follows from the fact that satisfying assignments are not always extendible. Indeed, consider two constraints $C_{i_1}, C_{i_2} \in L$ that share a common variable $j \in R$. Set $S_2 = T_{i_1} \cup T_{i_2}$, and $S_1 = S_2 \setminus \{j\}$.⁵ Assuming that the support of no other constraint is contained in S_2 , we get that distribution $\mathcal{D}(S_1)$ maps any variable in S_1 to $\{0, 1\}$ with probability $1/2$ independently, but some of these assignments are not even extendible to S_2 meaning that $\mathcal{D}(S_2)$ will assign them with probability zero.

Thus, to define $\mathcal{D}(S)$, we cannot simply sample assignments satisfying all constraints in $\mathcal{C}(S)$ with probabilities given by μ . In fact the above example shows that any attempt to blindly assign a set S with a distribution that is supported on all satisfying assignments for S is bound to fail. At the same time it seems hard to reason about a distribution that uses a totally different concept. To overcome this obstacle, we take a two step approach:

1. For a set S we define a superset \bar{S} such that \bar{S} is “global enough” to contain sufficient information, while it also is “local enough” so that $\mathcal{C}(\bar{S})$ is not too large. We require the property of such sets that if we remove \bar{S} and $\mathcal{C}(\bar{S})$, then the remaining graph $G|_{-\bar{S}}$ still has good expansion. We deal with this in Section 3.1.
2. When μ is the uniform distribution over $P^{-1}(1)$, the distribution $\mathcal{D}(S)$ is going to be the uniform distribution over satisfying assignments in \bar{S} . In the case that μ is not uniform over $P^{-1}(1)$, we give a natural generalization to the above uniformity. We show how to define distributions, which we denote by $\mathcal{P}_\mu(S)$, such that for $S_1 \subseteq S_2$, the distributions $\mathcal{P}_\mu(S_1)$ and $\mathcal{P}_\mu(S_2)$ are guaranteed to be consistent if $G|_{-S_1}$ has good expansion. This appears in Section 3.2.

We then combine the two techniques and define $\mathcal{D}(S)$ according to $\mathcal{P}_\mu(\bar{S})$. This is done in Section 4.

⁵Remember that T_{i_1} is the set of variables appearing in C_{i_1} and T_{i_2} is the set of variables appearing in C_{i_2} .

3.1 Finding advice-sets

We now give an algorithm below to obtain a superset \bar{S} for a given set S , which we call the advice-set of S . It is inspired by the “expansion correction” procedure in [8].

Algorithm Advice

The input is an (r, e_1) -boundary expanding bipartite graph $G = (L, R, E)$, some $e_2 \in (0, e_1)$, and some $S \subseteq R$, $|S| < (e_1 - e_2)r$, with some order $S = \{x_1, \dots, x_t\}$.

Initially set $\bar{S} \leftarrow \emptyset$ and $\xi \leftarrow r$

For $j = 1, \dots, |S|$ **do**

$M_j \leftarrow \emptyset$

$\bar{S} \leftarrow \bar{S} \cup \{x_j\}$

If $G|_{-\bar{S}}$ is not (ξ, e_2) -boundary expanding **then**

Find a maximal $M_j \subset L$ in $G|_{-\bar{S}}$, such that $|M_j| \leq \xi$ and $|\partial M_j| \leq e_2|M_j|$ in $G|_{-\bar{S}}$

$\bar{S} \leftarrow \bar{S} \cup \Gamma(M_j)$

$\xi \leftarrow \xi - |M_j|$

Return \bar{S}

Theorem 3.1. *Algorithm Advice, when ran with inputs G, e_1, e_2, r , and S , returns $\bar{S} \subseteq R$ such that*

(a) $G|_{-\bar{S}}$ is (ξ_S, e_2) -boundary expanding,

(b) $\xi_S \geq r - \frac{|S|}{e_1 - e_2}$, and

(c) $|\bar{S}| \leq \frac{(2e_1 + k - e_2)|S|}{2(e_1 - e_2)}$.

Proof. Let ξ_S be the value of ξ when the loop terminates. From the bounded size of M_j and how ξ changes at each iteration we know that ξ remains non-negative throughout the execution of the while loop, and in particular $\xi_S \geq 0$. Note that at step j , all the neighbors of M_j are added to the set \bar{S} so no member of M_j will be in $G_{-\bar{S}}$ after the j th step. In particular, all the sets M_j will be disjoint.

In order to prove (a) we will prove the following loop invariant: $G|_{-\bar{S}}$ is (ξ, e_2) -boundary expanding. Indeed, note that the input graph G is (ξ, e_1) -boundary expanding so the invariant holds at the beginning. At step j consider the set $\bar{S} \cup \{x_j\}$, and suppose that $G_{-(\bar{S} \cup \{x_j\})}$ is not (ξ, e_2) -boundary expanding. We find maximal M_j , $|M_j| \leq \xi$, such that $|\partial M_j| \leq e_2|M_j|$. We claim that $G_{-(\bar{S} \cup \{x_j\} \cup \Gamma(M_j))}$ is $(\xi - |M_j|, e_2)$ -boundary expanding. Assuming the contrary, there must be $M' \subset L$ such that $|M'| \leq \xi - |M_j|$ and $|\partial M'| \leq e_2|M'|$. As we mentioned above, M_j will be disjoint from the left vertices of $G_{-(\bar{S} \cup \{x_j\} \cup \Gamma(M_j))}$, and in particular it will be disjoint from M' . Consider then $M_j \cup M'$ and note that $|M_j \cup M'| \leq \xi$. More importantly (right before we added $\Gamma(M_j)$ to \bar{S}) $|\partial(M_j \cup M')| \leq e_2|M_j| + e_2|M'| = e_2|M_j \cup M'|$ contradicting the maximality of M_j ; (a) follows.

To show (b) we consider the set $M = \bigcup_{j=1}^t M_j$ and upperbound and lowerbound its boundary expansion in G in two different ways. Notice that as we mentioned M_j 's are disjoint, so $|M| = \sum_{j=1}^t |M_j| = r - \xi_S$. Since G is (r, e_1) -boundary expanding, the set M has at least $e_1(r - \xi_S)$ boundary neighbors in G . But each member of $\partial M \setminus S$ is in the boundary of exactly one of the M_j 's, so it will be counted towards the boundary expansion of M_j in $G_{-\bar{S}}$ in the j iteration of the loop (for some j). Given that M_j 's have boundary expansion at most e_2 we have

$$e_1(r - \xi_S) \leq |\partial M| \leq |S| + \sum_j e_2 |M_j| = |S| + e_2(r - \xi_S),$$

which readily implies (b).

Finally note that \bar{S} consists of S union the neighbors of all M_j 's. But given that M_j 's have boundary at most expansion e_2 they can not have a big neighbor set. In particular

$$|\bar{S}| \leq |S| + \sum_j \Gamma(M_j) \leq |S| + \frac{e_2 + k}{2} \sum_j |M_j| \leq |S| + \frac{(e_2 + k)|S|}{2(e_1 - e_2)} = \frac{(2e_1 + k - e_2)|S|}{2(e_1 - e_2)},$$

which proves (c). □

3.2 Defining the Distributions $\mathcal{P}_\mu(S)$

We now define for every set S , a distribution $\mathcal{P}_\mu(S)$ such that for any $\alpha \in [q]^S$, $\Pr_{\mathcal{P}_\mu(S)}[\alpha] > 0$ only if α satisfies all the constraints in $\mathcal{C}(S)$. For a constraint C_i with set of inputs T_i , defined as $C_i(x_{i_1}, \dots, x_{i_k}) \equiv P(x_{i_1} + a_{i_1}, \dots, x_{i_k} + a_{i_k})$, let $\mu_i : [q]^{T_i} \rightarrow [0, 1]$ denote the distribution

$$\mu_i(x_{i_1}, \dots, x_{i_k}) = \mu(x_{i_1} + a_{i_1}, \dots, x_{i_k} + a_{i_k})$$

so that the support of μ_i is contained in $C_i^{-1}(1)$. We then define the distribution $\mathcal{P}_\mu(S)$ by picking each assignment $\alpha \in [q]^S$ with probability proportional to $\prod_{C_i \in \mathcal{C}(S)} \mu_i(\alpha(T_i))$. Formally,

$$\Pr_{\mathcal{P}_\mu(S)}[\alpha] = \frac{1}{Z_S} \cdot \prod_{C_i \in \mathcal{C}(S)} \mu_i(\alpha(T_i)) \quad (3.1)$$

where $\alpha(T_i)$ is the restriction of α to T_i and Z_S is a normalization factor given by

$$Z_S = \sum_{\alpha \in [q]^S} \prod_{C_i \in \mathcal{C}(S)} \mu_i(\alpha(T_i)).$$

To understand the distribution, it is easier to think of the special case when μ is just the uniform distribution on $P^{-1}(1)$ (like in the case of MAX k-XOR). Then $\mathcal{P}_\mu(S)$ is simply the uniform distribution on assignments satisfying all the constraints in $\mathcal{C}(S)$. When μ is not uniform, the probabilities are weighted by the product of the values $\mu_i(\alpha(T_i))$ for all the constraints.⁶ However, we still have the property that if $\Pr_{\mathcal{P}_\mu(S)}[\alpha] > 0$, then α satisfies all the constraints in $\mathcal{C}(S)$.

In order for the distribution $\mathcal{P}_\mu(S)$ to be well defined, we need to ensure that $Z_S > 0$. The following lemma shows how to calculate Z_S if G is sufficiently expanding, and simultaneously proves that if $S_1 \subseteq S_2$, and if $G|_{-S_1}$ is sufficiently expanding, then $\mathcal{P}_\mu(S_1)$ is consistent with $\mathcal{P}_\mu(S_2)$ over S_1 .

⁶Note however that $\mathcal{P}_\mu(S)$ is not a product distribution because different constraints in $\mathcal{C}(S)$ may share variables.

Lemma 3.2. *Let Φ be a MAX k -CSP $_q(P)$ instance as above and $S_1 \subseteq S_2$ be two sets of variables such that both G and $G|_{-S_1}$ are $(r, k - 3 + \delta)$ -boundary expanding for some $\delta > 0$ and $|\mathcal{C}(S_2)| \leq r$. Then $Z_{S_2} = q^{|S_2|}/q^{k|\mathcal{C}(S_2)|}$, and for any $\alpha_1 \in [q]^{S_1}$*

$$\sum_{\substack{\alpha_2 \in [q]^{S_2} \\ \alpha_2(S_1) = \alpha_1}} \Pr_{\mathcal{P}_\mu(S_2)}[\alpha_2] = \Pr_{\mathcal{P}_\mu(S_1)}[\alpha_1]. \quad (3.2)$$

Proof. Let $\mathcal{C} = \mathcal{C}(S_2) \setminus \mathcal{C}(S_1)$ be the set of t constraints dominated by S_2 but not S_1 . Without loss of generality let $\mathcal{C} = \{C_1, \dots, C_t\}$ with C_i being on the set of variables T_i some of which might be set by α_1 . Note that any α_2 consistent with α_1 can be written as $\alpha_1 \circ \alpha$ for some $\alpha \in [q]^{S_2 \setminus S_1}$. We will show a way to calculate a sum similar to the left hand side of (3.2). Note that these calculations are meaningful even if Z_{S_1} or Z_{S_2} are zero, in which case both sides are simply zero. Taking $S_1 = \emptyset$ will then give us the value of Z_{S_2} .

$$\begin{aligned} Z_{S_2} \cdot \sum_{\substack{\alpha_2 \in [q]^{S_2} \\ \alpha_2(S_1) = \alpha_1}} \Pr_{\mathcal{P}_\mu(S_2)}[\alpha_2] &= \sum_{\alpha \in [q]^{S_2 \setminus S_1}} \prod_{C_i \in \mathcal{C}(S_2)} \mu_i((\alpha_1 \circ \alpha)(T_i)) \\ &= \left(\prod_{C_i \in \mathcal{C}(S_1)} \mu_i(\alpha_1(T_i)) \right) \sum_{\alpha \in [q]^{S_2 \setminus S_1}} \prod_{j=1}^t \mu_j((\alpha_1 \circ \alpha)(T_j)) \\ &= \left(Z_{S_1} \cdot \Pr_{\mathcal{P}_\mu(S_1)}[\alpha_1] \right) \sum_{\alpha \in [q]^{S_2 \setminus S_1}} \prod_{j=1}^t \mu_j((\alpha_1 \circ \alpha)(T_j)). \end{aligned}$$

The following claim lets us calculate this last sum conveniently using the expansion of $G|_{-S_1}$.

Claim 3.3. *Let \mathcal{C} be as above. Then there exists an ordering C_{i_1}, \dots, C_{i_t} of constraints in \mathcal{C} and a partition of $S_2 \setminus S_1$ into sets of variables F_1, \dots, F_t and F_{t+1} such that for all $j \leq t$, $F_j \subseteq T_{i_j}$, $|F_j| \geq k - 2$, and*

$$\forall j \ F_j \cap \left(\bigcup_{\ell > j} T_{i_\ell} \right) = \emptyset.$$

Proof of Claim 3.3. We build the sets F_j inductively using the fact that $G|_{-S_1}$ is $(r, k - 3 + \delta)$ -boundary expanding.

Start with the set of constraints $\mathcal{C}_1 = \mathcal{C}$. Since $|\mathcal{C}_1| = |\mathcal{C}(S_2) \setminus \mathcal{C}(S_1)| \leq r$, \mathcal{C}_1 should be expanding in $G|_{-S_1}$, i. e., $|\partial(\mathcal{C}_1) \setminus S_1| \geq (k - 3 + \delta)|\mathcal{C}_1|$. Hence, there exists $C_j \in \mathcal{C}_1$ contributing at least $k - 2$ variables to the boundary of \mathcal{C}_1 , i. e., $|T_j \cap (\partial(\mathcal{C}_1) \setminus S_1)| \geq k - 2$. Let $T_j \cap (\partial(\mathcal{C}_1) \setminus S_1) = F_1$ and $i_1 = j$. We then take $\mathcal{C}_2 = \mathcal{C}_1 \setminus \{C_{i_1}\}$ and continue in the same way. What is left from $S_2 \setminus S_1$ after the last step will be F_{t+1} . It is not hard to check that F_{t+1} is the portion of $S_2 \setminus S_1$ that is not in any \mathcal{C}_i although we will not use this.

Since at every step we have $F_j \subseteq \partial(\mathcal{C}_j) \setminus S_1$ and, for all $\ell > j$, $\mathcal{C}_\ell \subseteq \mathcal{C}_j$, F_j shares no variables with $\Gamma(\mathcal{C}_\ell)$ for $\ell > j$. Hence, we get $F_j \cap \left(\bigcup_{\ell > j} T_{i_\ell} \right) = \emptyset$ as claimed. \square

Using this decomposition, we can reorder the sum and split it as

$$\sum_{\alpha \in [q]^{S_2 \setminus S_1}} \prod_{j=1}^t \mu_j(\alpha_1 \circ \alpha(T_j)) = \sum_{\beta_{t+1} \in [q]^{F_{t+1}}} \sum_{\beta_t \in [q]^{F_t}} \mu_{i_t} \sum_{\beta_{t-1} \in [q]^{F_{t-1}}} \mu_{i_{t-1}} \cdots \sum_{\beta_2 \in [q]^{F_2}} \mu_{i_2} \sum_{\beta_1 \in [q]^{F_1}} \mu_{i_1}$$

where the input to each μ_j depends on α_1 and $\beta_j, \dots, \beta_{r+1}$ but not on $\beta_1, \dots, \beta_{j-1}$.

We now reduce the expression from right to left. Since F_1 contains at least $k-2$ variables and μ_{i_1} is a balanced pairwise independent distribution,

$$\sum_{\beta_1 \in [q]^{F_1}} \mu_{i_1} = \Pr_{\mu}[(\alpha_1 \circ \beta_2 \dots \circ \beta_r)(T_{i_1} \setminus F_1)] = \frac{1}{q^{k-|F_1|}}$$

irrespective of the values assigned by $\alpha_1 \circ \beta_2 \circ \dots \circ \beta_r$ to the remaining (at most 2) variables in $T_{i_1} \setminus F_1$. Continuing in this fashion from right to left, we get that

$$\sum_{\alpha \in [q]^{S_2 \setminus S_1}} \prod_{j=1}^t \mu_j((\alpha_1 \circ \alpha)(T_j)) = \frac{1}{q^{kt - \sum_{j=1}^{t+1} |F_j|}} = q^{|S_2 \setminus S_1| - k|\mathcal{C}(S_2) \setminus \mathcal{C}(S_1)|}.$$

Hence, we get that

$$Z_{S_2} \cdot \sum_{\substack{\alpha_2 \in [q]^{S_2} \\ \alpha_2(S_1) = \alpha_1}} \Pr_{\mathcal{P}_{\mu}(S_2)}[\alpha_2] = \left(Z_{S_1} \cdot \frac{q^{|S_2 \setminus S_1|}}{q^{k|\mathcal{C}(S_2) \setminus \mathcal{C}(S_1)|}} \right) \Pr_{\mathcal{P}_{\mu}(S_1)}[\alpha_1]. \quad (3.3)$$

Now, since we know that G is $(r, k-3+\delta)$ -boundary expanding, we can replace S_1 by \emptyset in the above calculation to get

$$Z_{S_2} = \sum_{\alpha \in [q]^{S_2}} \prod_{C_i \in \mathcal{C}(S_2)} \mu_i((\alpha_1 \circ \alpha)(T_i)) = q^{|S_2| - k|\mathcal{C}(S_2)|},$$

as we wanted. Plugging in the value of Z_{S_2} and Z_{S_1} into (3.3) will show that

$$\sum_{\substack{\alpha_2 \in [q]^{S_2} \\ \alpha_2(S_1) = \alpha_1}} \Pr_{\mathcal{P}_{\mu}(S_2)}[\alpha_2] = \Pr_{\mathcal{P}_{\mu}(S_1)}[\alpha_1],$$

which proves the lemma. \square

We are now almost ready to describe our Sherali-Adams SDP solution. The only other property of $\mathcal{P}_{\mu}(S)$ that we need to show is that it is pairwise-independent and balanced.

Claim 3.4. *Let G be a $(r, k-3+\varepsilon)$ -boundary expanding constraint graph, with $\varepsilon > 2/3$ in which no two constraints share more than one variable. Then for any $S \subset R$, $|S| \leq r$, the distribution $\mathcal{P}_{\mu}(S)$ is a pairwise-independent and balanced distribution. That is, for all $i_1, i_2 \in S$, $j_1, j_2 \in [q]$,*

$$\Pr_{\alpha \sim \mathcal{P}_{\mu}(S)}[\alpha_{i_1} = j_1] = 1/q, \quad \Pr_{\alpha \sim \mathcal{P}_{\mu}(S)}[(\alpha_{i_1} = j_1) \wedge (\alpha_{i_2} = j_2)] = 1/q^2.$$

Proof. Let S be a subset of the variables, $i, j \in S$ and $\varepsilon' = \min(1, \varepsilon - 2/3)$. We will first show that $G_{-\{i,j\}}$ is $(r, k-3+\varepsilon')$ -boundary expanding.

Let \mathcal{C} be a set of constraints with $|\mathcal{C}| \leq r$. When $|\mathcal{C}| = 1$, \mathcal{C} has k boundary neighbors in G and hence at least $k-2 \geq (k-3+\varepsilon')|\mathcal{C}|$ boundary neighbors in $G_{-\{i,j\}}$. When $|\mathcal{C}| = 2$, the number of boundary neighbors in G must be at least $2k-2$ since the two constraints in \mathcal{C} can share at most one variable. It follows that $|\partial \mathcal{C}|$ is at least $2k-2-2 \geq (k-3+\varepsilon')|\mathcal{C}|$ in $G_{-\{i,j\}}$.

Finally, when $3 \leq |\mathcal{C}| \leq r$, we get that the size of $|\partial\mathcal{C}|$ in $G|_{-\{i,j\}}$ is at least $(k-3+\varepsilon)|\mathcal{C}|-2$ as G is $(r, k-3+\varepsilon)$ -boundary expanding. This proves the claim as

$$(k-3+\varepsilon)|\mathcal{C}|-2 = (k-3+\varepsilon-2/|\mathcal{C}|)|\mathcal{C}| \geq (k-3+\varepsilon')|\mathcal{C}|.$$

It follows from applying [Lemma 3.3](#) with $S_2 = S$ and $S_1 = \{i, j\}$ that $\mathcal{P}_\mu(S)$ agrees with $\mathcal{P}_\mu(\{i, j\})$ on the set $\{i, j\}$. Now note that for $k=2$ the only promising predicate is the one accepting everything for which the theorem is trivial, so one can assume that $k > 2$ and $C(\{i, j\})$ is empty. It follows that $\mathcal{P}_\mu(\{i, j\})$ is the uniform distribution on $[q]^{\{i,j\}}$ which completes the proof. \square

4 Constructing the integrality gap

We now show how to construct integrality gaps using the ideas in the previous section. For a given promising predicate P , our integrality gap instance will be a random instance Φ of the MAX k -CSP $_q(P)$ problem, conditioned on no two constraints sharing more than one variable. To generate a random instance with m constraints, for every constraint C_i , we randomly select a k -tuple $T_i = \{x_{i_1}, \dots, x_{i_k}\}$ of distinct variables and $a_{i_1}, \dots, a_{i_k} \in [q]$, and put $C_i \equiv P(x_{i_1} + a_{i_1}, \dots, x_{i_k} + a_{i_k})$. It is well known and used in various works on integrality gaps and proof complexity (e. g., [\[8\]](#), [\[1, 2\]](#), [\[27\]](#) and [\[26\]](#)), that random instances of CSPs are both highly unsatisfiable and highly expanding. We capture the properties we need in the lemma below. The proof uses standard arguments; we defer it to the next section.

Lemma 4.1. *Let $\varepsilon, \delta > 0$ and a predicate $P : [q]^k \rightarrow \{0, 1\}$ be given. Then there exist $\gamma = O(q^k \log q / \varepsilon^2)$, $\eta = \Omega((1/\gamma)^{10/\delta})$ and $N \in \mathbb{N}$, such that if $n \geq N$ and Φ is a random instance of MAX k -CSP $_q(P)$ with $m = \gamma n$ constraints, then with probability $\exp(-O(k^4 \gamma^2))$*

1. $OPT(\Phi) \leq \frac{|P^{-1}(1)|}{q^k} (1 + \varepsilon) \cdot m$,
2. for any set \mathcal{C} of constraints with $|\mathcal{C}| \leq \eta n$, we have $|\partial(\mathcal{C})| \geq (k-2-\delta)|\mathcal{C}|$, and
3. no two constraints in Φ share more than one variable.

Let Φ be an instance of MAX k -CSP $_q$ on n variables for which G_Φ is $(\eta n, k-2-\delta)$ -boundary expanding for some $\delta < 1/4$, as in [Lemma 4.1](#). For such a Φ , we now define the distributions $\mathcal{D}(S)$. In the rest of this chapter we assume $k \geq 3$ as for $k=2$ the only promising predicate is the one satisfying all assignments for which the theorem is trivial.

For a set S of size at most $t = \eta n / 6k$, let \bar{S} be subset of variables output by the algorithm Advice when run with input S and parameters $r = \eta n$, $e_1 = (k-2-\delta)$, $e_2 = (k-8/3-\delta)$ on the graph G_Φ . [Theorem 3.1](#) shows that

$$|\bar{S}| \leq \frac{(2k-4-2\delta+k-k+8/3+\delta)|S|}{4/3} = \frac{(6k-4-3\delta)|S|}{4} \leq \frac{\eta n}{4},$$

and, also,

$$\xi_S \geq \eta n - \frac{|S|}{2/3} = \eta n - \frac{\eta n}{4k} > \frac{3\eta n}{4}.$$

We then use (3.1) to define the distribution $\mathcal{D}(S)$ for sets S of size at most $\eta n/6k$ as

$$\Pr_{\mathcal{D}(S)}[\alpha] = \sum_{\substack{\beta \in [q]^{\bar{S}} \\ \beta(S)=\alpha}} \Pr_{\mathcal{P}_\mu(\bar{S})}[\beta].$$

Using the properties of the distributions $\mathcal{P}_\mu(\bar{S})$, we can now prove that the distributions $\mathcal{D}(S)$ are consistent.

Claim 4.2. *Let the distributions $\mathcal{D}(S)$ be defined as above. Then for any two sets $S_1 \subseteq S_2 \subseteq [n]$ with $|S_2| \leq t = \eta n/6k$, the distributions $\mathcal{D}(S_1), \mathcal{D}(S_2)$ are equal on S_1 .*

Proof. The distributions $\mathcal{D}(S_1)$ and $\mathcal{D}(S_2)$ are defined according to $\mathcal{P}_\mu(\bar{S}_1)$ and $\mathcal{P}_\mu(\bar{S}_2)$, respectively. To prove the claim, we show that $\mathcal{P}_\mu(\bar{S}_1)$ and $\mathcal{P}_\mu(\bar{S}_2)$ are equal to the distribution $\mathcal{P}_\mu(\bar{S}_1 \cup \bar{S}_2)$ on \bar{S}_1 and \bar{S}_2 , respectively (note that it need not be the case that $\bar{S}_1 \subseteq \bar{S}_2$).

Let $S_3 = \bar{S}_1 \cup \bar{S}_2$. Since $|\bar{S}_1|, |\bar{S}_2| \leq \eta n/4$, we have $|S_3| \leq \eta n/2$. We will first show that $|\mathcal{C}(S_3)| \leq 3\eta n/4$. Assume to the contrary, and let \mathcal{C} be any subset of $\mathcal{C}(S_3)$ of size $3\eta n/4$. Given that $k \geq 3$ and $\delta < 1/4$ we would have the following bound on the boundary of \mathcal{C} ,

$$\frac{|\partial \mathcal{C}|}{|\mathcal{C}|} \leq \frac{|\Gamma(\mathcal{C})|}{|\mathcal{C}|} \leq \frac{|S_3|}{|\mathcal{C}|} \leq \frac{1/2}{3/4} = \frac{2}{3} < k - 2 - \delta,$$

which would contradict boundary expansion of G .

Now, by Theorem 3.1, we know that both $G|_{-\bar{S}_1}$ and $G|_{-\bar{S}_2}$ are $(3\eta n/4, k - 8/3 - \delta)$ -boundary expanding. Thus, using Lemma 3.2 for the pairs $\bar{S}_1 \subseteq S_3$ and $\bar{S}_2 \subseteq S_3$, we get that

$$\begin{aligned} \Pr_{\mathcal{D}(S_1)}[\alpha_1] &= \sum_{\substack{\beta_1 \in [q]^{\bar{S}_1} \\ \beta_1(S_1)=\alpha_1}} \Pr_{\mathcal{P}_\mu(\bar{S}_1)}[\beta_1] = \sum_{\substack{\beta_3 \in [q]^{\bar{S}_3} \\ \beta_3(S_1)=\alpha_1}} \Pr_{\mathcal{P}_\mu(S_3)}[\beta_3] \\ &= \sum_{\substack{\beta_2 \in [q]^{\bar{S}_2} \\ \beta_2(S_1)=\alpha_1}} \Pr_{\mathcal{P}_\mu(\bar{S}_2)}[\beta_2] = \sum_{\substack{\alpha_2 \in [q]^{\bar{S}_2} \\ \alpha_2(S_1)=\alpha_1}} \Pr_{\mathcal{D}(S_2)}[\alpha_2] \end{aligned}$$

which shows that $\mathcal{D}(S_1)$ and $\mathcal{D}(S_2)$ are equal on S_1 . \square

It is now easy to prove the main result.

Theorem 4.3. *Let $P : [q]^k \rightarrow \{0, 1\}$ be a promising predicate. Then for every constant $\zeta > 0$, there exist $c = c(q, k, \zeta)$, such that for large enough n , the integrality gap of MAX k -CSP $_q(P)$ for the program obtained by cn levels of the Sherali-Adams SDP hierarchy is at least*

$$\frac{q^k}{|P^{-1}(1)|} - \zeta.$$

Proof. We take $\varepsilon = \zeta/q^k$, $\delta = 1/4$ and consider a random instance Φ of MAX k -CSP $_q(P)$ with $m = \gamma n$ as given by Lemma 4.1. Thus, $\text{OPT}(\Phi) \leq \frac{|P^{-1}(1)|}{q^k} (1 + \varepsilon) \cdot m$.

On the other hand, by [Claim 4.2](#) we can define distributions $\mathcal{D}(S)$ over every set of at most $\eta n/6k$ variables such that for $S_1 \subseteq S_2$, $\mathcal{D}(S_1)$ and $\mathcal{D}(S_2)$ are consistent over S_1 . Also, [Claim 3.4](#) implies that $\mathcal{P}_\mu(\bar{S})$ hence $\mathcal{D}(S)$ is pairwise-independent and balanced for any S . We can now construct the SDP vectors with inner products agreeing with the probabilities according to the distributions $\mathcal{D}(S) = \mathcal{P}_\mu(\bar{S})$ for an instance satisfying the above property using the following simple fact.

Claim 4.4. *There exist vectors $\{\mathbf{v}_{(i,j)}\}_{i \in [n], j \in [q]}$ and \mathbf{v}_0 satisfying:*

1. for all $i \in [n]$ and $j_1 \neq j_2$, $\langle \mathbf{v}_{(i,j_1)}, \mathbf{v}_{(i,j_2)} \rangle = 0$,
2. for all $i \in [n], j \in [q]$, $\langle \mathbf{v}_{(i,j)}, \mathbf{v}_0 \rangle = \|\mathbf{v}_{(i,j)}\|^2 = 1/q$, and
3. for all $i_1 \neq i_2 \in [n]$ and $j_1, j_2 \in [q]$, $\langle \mathbf{v}_{(i_1,j_1)}, \mathbf{v}_{(i_2,j_2)} \rangle = 1/q^2$.

Proof. Let $\mathbf{e}_1, \dots, \mathbf{e}_n$ be an orthonormal basis for \mathbb{R}^n and $\mathbf{u}_0, \dots, \mathbf{u}_{q-1}$ be vertices of a $q-1$ dimensional simplex satisfying $\langle \mathbf{u}_{j_1}, \mathbf{u}_{j_2} \rangle = -1/(q-1)$ when $j_1 \neq j_2$ and 1 otherwise. Let $\mathbf{v}_0 \in \mathbb{R}^{n(q-1)+1}$ be a unit vector such that $\langle \mathbf{v}_0, \mathbf{e}_i \otimes \mathbf{u}_j \rangle = 0$ for all $i \in [n]$ and $j \in [q]$. We then define $\mathbf{v}_{(i,j)}$ as

$$\mathbf{v}_{(i,j)} := \frac{1}{q} \mathbf{v}_0 + \frac{\sqrt{q-1}}{q} (\mathbf{e}_i \otimes \mathbf{u}_j).$$

It is easy to check that $\langle \mathbf{v}_0, \mathbf{v}_{(i,j_1)} \rangle = 1/q$ and $\langle \mathbf{v}_{(i_1,j_1)}, \mathbf{v}_{(i_2,j_2)} \rangle = 1/q^2$ for all $i_1 \neq i_2 \in [n]$ and $j_1, j_2 \in [q]$. Also, for $j_1 \neq j_2$

$$\langle \mathbf{v}_{(i,j_1)}, \mathbf{v}_{(i,j_2)} \rangle = \frac{1}{q^2} + \frac{q-1}{q^2} \cdot \left(\frac{-1}{q-1} \right) = 0$$

which proves the claim. □

By [Lemma 2.3](#) this gives a feasible solution to the SDP obtained by $\eta n/(6k)$ levels. Also, by definition of $\mathcal{D}(S)$, we have that $\Pr_{\mathcal{D}(S)}[\alpha] > 0$ only if α satisfies all constraints in $\mathcal{C}(\bar{S}) \supseteq \mathcal{C}(S)$. Hence, the value of $\text{FRAC}(\Phi)$ is given by

$$\sum_{i=1}^m \sum_{\alpha \in [q]^{T_i}} C_i(\alpha) X_{(T_i, \alpha)} = \sum_{i=1}^m \sum_{\alpha \in [q]^{T_i}} C_i(\alpha) \Pr_{\mathcal{D}(T_i)}[\alpha] = \sum_{i=1}^m \sum_{\alpha \in [q]^{T_i}} \Pr_{\mathcal{D}(T_i)}[\alpha] = m.$$

Thus, the integrality gap after $\eta n/(6k)$ levels is at least

$$\frac{\text{FRAC}(\Phi)}{\text{OPT}(\Phi)} = \frac{q^k}{|P^{-1}(1)|(1+\epsilon)} \geq \frac{q^k}{|P^{-1}(1)|} - \zeta.$$

□

5 Proof of Lemma 4.1

The proof is a simple modification of the proof of [27, Lemma 5]. We include it for convenience.

Proof. Let $\alpha \in [q]^n$ be any fixed assignment. For a fixed α , the events that a constraint C_i is satisfied are independent and happen with probability $|P^{-1}(1)|/q^k$ each. Hence, the probability over the choice of Φ that α satisfies more than $(|P^{-1}(1)|/q^k)(1 + \varepsilon) \cdot \gamma n$ constraints is at most

$$\exp\left(-\frac{\varepsilon^2 \gamma n |P^{-1}(1)|}{3q^k}\right)$$

by Chernoff bounds. By a union bound, the probability that *any* assignment satisfies more than $(|P^{-1}(1)|/q^k)(1 + \varepsilon) \cdot \gamma n$ constraints is at most

$$q^n \cdot \exp\left(-\frac{\varepsilon^2 \gamma n |P^{-1}(1)|}{3q^k}\right) = \exp\left(n \ln q - \frac{\varepsilon^2 \gamma n |P^{-1}(1)|}{3q^k}\right)$$

which is $o(1)$ for $\gamma = (6q^k \ln q)/\varepsilon^2$.

For showing the boundary expansion, we note that it suffices to show that the constraints have large expansion, i. e., each set of s constraints (for $s \leq \eta n$) contains at least $(k - 1 - \delta/2)s$ variables. Since each non-boundary variable occurs in at least two constraints, we get that the number of boundary variables must be at least $2(k - 1 - \delta/2)s - ks = (k - 2 - \delta)s$.

To show this, consider the probability that a set of s constraints contains at most cs variables, where $c = k - 1 - \delta$. This is upper bounded by

$$\binom{n}{cs} \cdot \binom{cs}{s} \cdot s! \binom{\gamma n}{s} \cdot \binom{n}{k}^{-s}.$$

Here $\binom{n}{cs}$ is the number of ways of choosing the cs variables involved, $\binom{cs}{s}$ is the number of ways of picking s tuples out of all possible k -tuples on cs variables and $s! \binom{\gamma n}{s}$ is the number of ways of selecting the s constraints. The number $\binom{n}{k}^s$ is simply the number of ways of picking s of these k -tuples in an unconstrained way. Using $\left(\frac{a}{b}\right)^b \leq \frac{a}{b} \leq \left(\frac{a-e}{b}\right)^b$, $s! \leq s^s$ and collecting terms, we can bound this expression by

$$\left(\frac{s}{n}\right)^{\delta s/2} \left(e^{2k+1-\delta/2} k^{1+\delta/2} \gamma\right)^s \leq \left(\frac{s}{n}\right)^{\delta s/2} (\gamma^5)^s = \left(\frac{s \gamma^{10/\delta}}{n}\right)^{\delta s/2}.$$

We need to show that the probability that a set of s constraints contains less than cs variables for *any* $s \leq \eta n$ is $o(1)$. Thus, we sum this probability over all $s \leq \eta n$ to get

$$\begin{aligned} \sum_{s=1}^{\eta n} \left(\frac{s \gamma^{10/\delta}}{n}\right)^{\delta s/2} &= \sum_{s=1}^{\ln^2 n} \left(\frac{s \gamma^{10/\delta}}{n}\right)^{\delta s/2} + \sum_{s=\ln^2 n+1}^{\eta n} \left(\frac{s \gamma^{10/\delta}}{n}\right)^{\delta s/2} \\ &\leq O\left(\frac{\gamma^{10}}{n^{\delta/2}} \ln^2 n\right) + O\left(\left(\eta \cdot \gamma^{10/\delta}\right)^{(\delta/2) \ln^2 n}\right). \end{aligned}$$

The first term is $o(1)$ and is small for large n . The second term is also $o(1)$ for $\eta = 1/(100\gamma^{10/\delta})$.

Thus, we get the first two properties with probability $1 - o(1)$. Finally, the probability that there are no two constraints sharing two variables must be at least $\prod_{i=1, \dots, m} (1 - O(i \cdot k^4/n^2))$ because when we choose the i th constraint, by wanting it to not share two variables with another previously chosen constraint, we are forbidding any of the $\binom{k}{2}$ pairs of variables in the i th constraint from being equal to any of the $(i-1) \cdot \binom{k}{2}$ pairs in the previously chosen ones. Now using that for small enough x , $1 - x > \exp(-2x)$, the probability is at least

$$\exp\left(-O\left(\frac{\sum_{i=1}^m i \cdot k^4}{n^2}\right)\right) = \exp\left(-O\left(\frac{k^4 \cdot m^2}{n^2}\right)\right) = \exp(-O(k^4 \gamma^2)).$$

□

Acknowledgements

The authors would like to thank the anonymous reviewers for suggestions that improved the presentation of the paper.

References

- [1] MIKHAIL ALEKHNovich, SANJEEV ARORA, AND IANNIS TOURLAKIS: Towards strong nonapproximability results in the Lovász-Schrijver hierarchy. In *Proc. 37th STOC*, pp. 294–303. ACM Press, 2005. [doi:10.1145/1060590.1060634] 271, 281
- [2] MIKHAIL ALEKHNovich, SANJEEV ARORA, AND IANNIS TOURLAKIS: Towards strong nonapproximability results in the Lovász-Schrijver hierarchy. *Comput. Complexity*, 20(4):615–648, 2011. [doi:10.1007/s00037-011-0027-z] 271, 281
- [3] SANJEEV ARORA, BÉLA BOLLOBÁS, LÁSZLÓ LOVÁSZ, AND IANNIS TOURLAKIS: Proving integrality gaps without knowing the linear program. *Theory of Computing*, 2(1):19–51, 2006. Preliminary version in *FOCS'02*. [doi:10.4086/toc.2006.v002a002] 271
- [4] PER AUSTRIN AND JOHAN HÅSTAD: Randomly supported independence and resistance. *SIAM J. Comput.*, 40(1):1–27, 2011. Preliminary version in *STOC'09*. [doi:10.1137/100783534] 270
- [5] PER AUSTRIN AND ELCHANAN MOSSEL: Approximation resistant predicates from pairwise independence. *Comput. Complexity*, 18(2):249–271, 2009. Preliminary version in *CCC'08*. [doi:10.1007/s00037-009-0272-6] 270, 276
- [6] MOHAMMAD HOSSEIN BATENI, MOSES CHARIKAR, AND VENKATESAN GURUSWAMI: MaxMin allocation via degree lower-bounded arborescences. In *Proc. 41st STOC*, pp. 543–552. ACM Press, 2009. [doi:10.1145/1536414.1536488] 271

- [7] ADITYA BHASKARA, MOSES CHARIKAR, EDEN CHLAMTAC, URIEL FEIGE, AND ARAVINDAN VIJAYARAGHAVAN: Detecting high log-densities—an $O(n^{1/4})$ approximation for densest k -subgraph. In *Proc. 42nd STOC*, pp. 201–210. ACM Press, 2010. [[doi:10.1145/1806689.1806719](https://doi.org/10.1145/1806689.1806719)] [271](#)
- [8] JOSHUA BURESH-OPPENHEIM, NICOLA GALESÌ, SHLOMO HOORY, AVNER MAGEN, AND TONIANN PITASSI: Rank bounds and integrality gaps for cutting planes procedures. *Theory of Computing*, 2(4):65–90, 2006. Preliminary version in *FOCS'03*. [[doi:10.4086/toc.2006.v002a004](https://doi.org/10.4086/toc.2006.v002a004)] [271](#), [277](#), [281](#)
- [9] MOSES CHARIKAR, KONSTANTIN MAKARYCHEV, AND YURY MAKARYCHEV: Integrality gaps for Sherali–Adams relaxations. In *Proc. 41st STOC*, pp. 283–292. ACM Press, 2009. [[doi:10.1145/1536414.1536455](https://doi.org/10.1145/1536414.1536455)] [272](#)
- [10] EDEN CHLAMTAC: Approximation algorithms using hierarchies of semidefinite programming relaxations. In *Proc. 48th FOCS*, pp. 691–701. IEEE Comp. Soc. Press, 2007. [[doi:10.1109/FOCS.2007.72](https://doi.org/10.1109/FOCS.2007.72)] [271](#)
- [11] EDEN CHLAMTAC AND GYANIT SINGH: Improved approximation guarantees through higher levels of SDP hierarchies. In *Proc. 11th Internat. Workshop on Approximation Algorithms for Combinatorial Optimization Problems (APPROX'08)*, volume 5171 of *Lecture Notes in Computer Science*, pp. 49–62. Springer, 2008. [[doi:10.1007/978-3-540-85363-3_5](https://doi.org/10.1007/978-3-540-85363-3_5)] [271](#)
- [12] LARS ENGBRETSSEN AND JONAS HOLMERIN: More efficient queries in PCPs for NP and improved approximation hardness of maximum CSP. *Random Structures Algorithms*, 33:497–514, 2008. Preliminary version in *STACS'05*. [[doi:10.1002/rsa.20226](https://doi.org/10.1002/rsa.20226)] [270](#)
- [13] URIEL FEIGE AND ROBERT KRAUTHGAMER: The probable value of the Lovász-Schrijver relaxations for maximum independent set. *SIAM J. Comput.*, 32(2):345–370, 2003. [[doi:10.1137/S009753970240118X](https://doi.org/10.1137/S009753970240118X)] [271](#)
- [14] WENCESLAS FERNANDEZ DE LA VEGA AND CLAIRE KENYON-MATHIEU: Linear programming relaxations of maxcut. In *Proc. 18th Ann. ACM-SIAM Symp. on Discrete Algorithms (SODA'07)*, pp. 53–61. SIAM, 2007. [[ACM:1283390](https://arxiv.org/abs/1283390)] [271](#)
- [15] KONSTANTINOS GEORGIU, AVNER MAGEN, TONIANN PITASSI, AND IANNIS TOURLAKIS: Integrality gaps of $2 - o(1)$ for Vertex Cover SDPs in the Lovász-Schrijver hierarchy. *SIAM J. Comput.*, 39(8):3553–3570, 2010. Preliminary version in *FOCS'07*. [[doi:10.1137/080721479](https://doi.org/10.1137/080721479)] [271](#)
- [16] VENKATESAN GURUSWAMI AND PRASAD RAGHAVENDRA: Constraint satisfaction over a non-Boolean domain: Approximation algorithms and Unique-Games hardness. In *Proc. 11th Internat. Workshop on Approximation Algorithms for Combinatorial Optimization Problems (APPROX'08)*, pp. 77–90. Springer, 2008. [[doi:10.1007/978-3-540-85363-3_7](https://doi.org/10.1007/978-3-540-85363-3_7)] [270](#)
- [17] JEAN B. LASSERRE: An explicit equivalent positive semidefinite program for nonlinear 0-1 programs. *SIAM J. Optim.*, 12(3):756–769, 2002. [[doi:10.1137/S1052623400380079](https://doi.org/10.1137/S1052623400380079)] [271](#)

- [18] MONIQUE LAURENT: A comparison of the Sherali-Adams, Lovász-Schrijver, and Lasserre relaxations for 0-1 programming. *Math. Oper. Res.*, 28(3):470–496, August 2003. [doi:10.1287/moor.28.3.470.16391] 271, 274
- [19] L. LOVÁSZ AND A. SCHRIJVER: Cones of matrices and set-functions and 0-1 optimization. *SIAM J. Optim.*, 1(2):166–190, May 1991. [doi:10.1137/0801013] 271
- [20] AVNER MAGEN AND MOHAMMAD MOHARRAMI: Robust algorithms for MAX INDEPENDENT SET on minor-free graphs based on the Sherali-Adams hierarchy. In *Proc. 12th Internat. Workshop on Approximation Algorithms for Combinatorial Optimization Problems (APPROX'09)*, pp. 258–271, Berlin, Heidelberg, 2009. Springer-Verlag. [doi:10.1007/978-3-642-03685-9_20] 271
- [21] CLAIRE MATHIEU AND ALISTAIR SINCLAIR: Sherali-Adams relaxations of the matching polytope. In *Proc. 41st STOC*, pp. 293–302. ACM Press, 2009. [doi:10.1145/1536414.1536456] 271
- [22] PRASAD RAGHAVENDRA: Optimal algorithms and inapproximability results for every CSP? In *Proc. 40th STOC*, pp. 245–254. ACM Press, 2008. [doi:10.1145/1374376.1374414] 271, 272
- [23] PRASAD RAGHAVENDRA AND DAVID STEURER: Integrality gaps for strong SDP relaxations of unique games. In *Proc. 50th FOCS*, pp. 575–585. IEEE Comp. Soc. Press, 2009. [doi:10.1109/FOCS.2009.73] 272
- [24] ALEX SAMORODNITSKY AND LUCA TREVISAN: Gowers uniformity, influence of variable, and PCPs. *SIAM J. Comput.*, 39:323–360, 2009. Preliminary version in *STOC'06*. [doi:10.1137/070681612] 270
- [25] ALEX SAMORODNITSKY AND LUCA TREVISAN: A PCP characterization of NP with optimal amortized query complexity. In *Proc. 32nd STOC*, pp. 191–199. ACM Press, 2010. [doi:10.1145/335305.335329] 270
- [26] GRANT SCHOENEBECK: Linear level Lasserre lower bounds for certain k -CSPs. In *Proc. 49th FOCS*, pp. 593–602. IEEE Comp. Soc. Press, 2008. [doi:10.1109/FOCS.2008.74] 271, 281
- [27] GRANT SCHOENEBECK, LUCA TREVISAN, AND MADHUR TULSIANI: A linear round lower bound for Lovász-Schrijver SDP relaxations of vertex cover. In *Proc. 22nd IEEE Conf. on Computational Complexity (CCC'07)*, pp. 205–216. IEEE Comp. Soc. Press, 2007. [doi:10.1109/CCC.2007.2] 271, 281, 284
- [28] GRANT SCHOENEBECK, LUCA TREVISAN, AND MADHUR TULSIANI: Tight integrality gaps for Lovász-Schrijver LP relaxations of vertex cover and max cut. In *Proc. 39th STOC*, pp. 302–310. ACM Press, 2007. [doi:10.1145/1250790.1250836] 271
- [29] HANIF D. SHERALI AND WARREN P. ADAMS: A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. *SIAM J. Discrete Math.*, 3(3):411–430, August 1990. [doi:10.1137/0403036] 271, 274

- [30] IANNIS TOURLAKIS: Towards optimal integrality gaps for hypergraph vertex cover in the Lovász-Schrijver hierarchy. In *Proc. 8th Internat. Workshop on Approximation Algorithms for Combinatorial Optimization Problems (APPROX'05)*, volume 3624, pp. 233–244. Springer, 2005. [[doi:10.1007/11538462_20](https://doi.org/10.1007/11538462_20)] 271
- [31] IANNIS TOURLAKIS: New lower bounds for vertex cover in the Lovász-Schrijver hierarchy. In *Proc. 21st IEEE Conf. on Computational Complexity (CCC'06)*, pp. 170–182. IEEE Comp. Soc. Press, 2006. [[doi:10.1109/CCC.2006.28](https://doi.org/10.1109/CCC.2006.28)] 271
- [32] MADHUR TULSIANI: CSP gaps and reductions in the Lasserre hierarchy. In *Proc. 41st STOC*, pp. 303–312. ACM Press, 2009. [[doi:10.1145/1536414.1536457](https://doi.org/10.1145/1536414.1536457)] 271

AUTHORS

Siavosh Benabbas [[About the author](#)]

PhD Candidate

University of Toronto, Toronto, Ontario, Canada

siavosh@cs.toronto.edu

<http://www.cs.toronto.edu/~siavosh>

Konstantinos Georgiou [[About the author](#)]

Post-doctoral fellow

Department of Combinatorics and Optimization

Faculty of Mathematics

University of Waterloo, Waterloo, Ontario, Canada

k2georgiou@math.uwaterloo.ca

<http://www.math.uwaterloo.ca/~k2georgi/>

Avner Magen [[About the author](#)]

Formerly Associate Professor

University of Toronto, Toronto, Ontario, Canada

<http://www.cs.toronto.edu/~avner>

Madhur Tulsiani [[About the author](#)]

Assistant Professor

Toyota Technological Institute at Chicago, Chicago, IL, USA

madhurt@ttic.edu

<http://ttic.uchicago.edu/~madhurt/>

ABOUT THE AUTHORS

SIAVOSH BENABBAS got his B. S. from [Sharif University of Technology](#) under the supervision of [Mohammad Ghodsi](#) in the [Department of Computer Engineering](#). Since 2006, Siavosh has been a graduate student in the [Department of Computer Science](#) at the [University of Toronto](#). There, he received his M. S. under the supervision of [Avner Magen](#) and [Charles Rackoff](#), while he is currently anticipating his graduation with a Ph. D. supervised by [Toni Pitassi](#). Siavosh is interested in questions touching on Linear Programming and Semidefinite Programming Relaxations, their Integrality Gaps and their applications to Combinatorial Optimization.

KONSTANTINOS GEORGIU (or Costis as friends call him) did his undergraduate studies at the [Department of Mathematics](#) at the [University of Athens](#) (1998-2002), from where he also received his M. S. in Logic and Algorithms (2002-2004). In 2005 he moved to Toronto where he received his Ph. D. under the supervision of [Avner Magen](#) and [Toni Pitassi](#) (2010). Currently, he is a postdoctoral fellow at the [Department of Combinatorics and Optimization](#) at the [University of Waterloo](#), working on projects varying from Convex Optimization to Complexity Theory to Combinatorial Optimization to Algorithmic Game Theory. Costis spends his free time in his kitchen where he experiments with flavors and recipes.

AVNER MAGEN (1968-2010) did his undergraduate and graduate studies at [Hebrew University](#) and received his Ph. D. in [Computer Science](#) in 2002 under the supervision of [Nati Linial](#). After holding a postdoctoral fellowship at [NEC Research](#) in Princeton, NJ, he joined the [University of Toronto](#) in 2002, first as a postdoctoral fellow, and then as an Assistant Professor in 2004. He was promoted to Associate Professor in 2009. Avner was a superb scholar, making fundamental contributions to a number of areas of theoretical computer science that include Metric Embeddings, Sublinear Algorithms, Convex Programming, Computational Geometry, and Approximation Algorithms. He was a wonderful colleague with a terrific sense of humor and great energy. He was a dedicated research supervisor and a superb teacher. He died at 42 in an avalanche while on a climbing tour in Alaska. He is sorely missed by his family, friends, colleagues, and students.

MADHUR TULSIANI did his B. Tech. in [Computer Science](#) at [IIT Kanpur](#), mentored by Professor [Somenath Biswas](#) (2001-05). He completed his Ph. D. at [UC Berkeley](#) (2005-09) under [Luca Trevisan](#). After graduation he spent two years as a postdoc at the [Institute for Advanced Study and Princeton University](#), and is currently an assistant professor at the [Toyota Technological Institute at Chicago](#). His research interests include Complexity Theory, Optimization, Approximation and Inapproximability, Pseudorandomness, and Cryptography.